









## IT HAPPENED IN NEW ENGLAND

News of General Interest  
From the Six States

Expenditures of the State of Connecticut during the 1923 fiscal year totaled \$21,172,846, or \$14.47 per capita, figures made public by the census bureau revealed.

Dr. Harold W. Hersey, superintendent of Bridgeport, Ct., Hospital, announced that a person whose name is withheld, has given the hospital \$25,000 for the purchase of 460 additional milligrams of radium.

The 19th annual farmers' week at the College of Agriculture, Orono, Me., will be held March 31, April 1 and 2. This will be the largest gathering of Maine farmers and homemakers ever attempted.

Sixteen prize Holstein cows, valued at \$150 each, were burned to death in a spectacular fire which destroyed a large barn on the estate of William Kings, Swampscott, Mass. Ten tons of hay were destroyed.

After falling 50 feet from the tower of a building under construction in Danvers, Mass., Fred Soucy, 40, picked himself up and drove an automobile 23 miles to his home in Methuen.

Horace S. Aldrich, formerly one of the wealthiest men in the State, died at his home in Milford, Mass. Mr. Aldrich, who was 75 years of age, weighed more than 500 pounds in his earlier years, but for some time had not weighed more than 300 pounds.

The recent earthquake was responsible for a decided decrease in the number of eggs collected at the chicken farm of James Ahern in Williamstown, Mass. Instead of the customary 75 or 80, less than a dozen were found when he made his daily visit to the henhouse.

The right arm of Brounau Bon-nassage was torn from his body while he was at work on a shafting in the bleaching department of the York Manufacturing Company, Saco, Me. His clothing caught in the shafting and caused the accident, which also resulted in three broken ribs and internal injuries.

President Coolidge has been invited by Governor Billings of Vermont, his native state, to attend an old home celebration to be held during the summer at any time and place in the state acceptable to the chief executive. Mr. Coolidge said he hoped he would be able to participate in the celebration.

Joseph H. Mahoney and Abel Friastron of Everett, Mass., in City Court, Montpelier, Vt., changed their plea of not guilty to the charge of transportation of liquor, admitting the guilt, and were fined \$1000 each. They were arrested with nine 1500 quarts of intoxicating liquor in their possession.

Miss Lizzie M. Cole, 62, died as she was preparing breakfast for her brother in the kitchen of their home in Boston, Mass. Heart trouble brought about the end as suddenly as it came, and in death she remained erect, unsupported, in the center of the floor, where one sister had been halted before she could lift her foot for another.

The committee on State administration reported the bill of the United Improvement Association for the continuance of the Massachusetts special commission on necessities of life. The committee recommends that the life of the State Commission on Necessaries of Life be extended until 1927 instead of 1927, as advocated by the United Improvement Association.

Judge Morton of the Superior Court, Boston, in a decree, declared Philip Berninger of Melrose Highlands is totally incapacitated for further work from inhalation of emery and metal dust while engaged as an operative and foreman in the Chelsea Clock Company plant, and sustained the original award of \$115 a week granted him by a single member of the industrial accident board.

Mrs. Chester Colson and her son, Alonzo Crawford, both of Bangor, are among 40 persons, most of whom live in Canada, who lay claim to a share of the \$7,000,000 Crawford estate now the subject of litigation in New York courts. The history of the Crawford estate goes back more than a century. It was accumulated by Andrew Crawford, who emigrated to Ontario, Canada, many years ago. In three sons, one, George, was the only one to marry, and it is through relationship with him that the Bangor woman and her son have their claim to the fortune, which is estimated at \$7,000,000 and is divided into three parts, securities and cash of the land.

Mrs. Amelia Hawsey, 92 years of age, who told a playful story of being snatched from the home of one relative to another when found without resources at the Attleboro, Mass., railroad station last July, was forced to make one more move when she was taken from the city home to the State Home at Tewksbury. Her son, George Hawsey, is serving a term in the House of Correction because he refused to provide for his aged mother. He is said to own real estate in Attleboro and in Central Falls, R. I.

## BOSTON MARKET REVIEW

Prepared by the Boston Office of the Bureau of Agricultural Economics, U. S. Department of Agriculture  
For Week Ending March 7, 1925

**FRUITS AND VEGETABLES:** Prices declined for the most part during the week, principally on account of heavy receipts. Spinach and cabbage declined noticeably. Maine potatoes were slightly weaker, closing at 1.20 per 100 lb. sack of Green Mt. Apples were about unchanged in price, but demand was slow, and considerable soft stock was on the market. Onions continued dull and weak, with most 100 lb. sacks of N. Y. and Maine Yellow Globes selling around 2.50-2.60. Pickles were weaker at 2.00-2.10 per 100 lb. sack. Chile Valley shipping points were slightly advanced, principally on account of the advance in the New York City market. Shipments were light with sales ranging from 250-300 tons. Potatoes were lower at 2.00-2.10. Beans were lower at 2.00-2.10. Peas were lower at 2.00-2.10. Corn was lower at 2.00-2.10. Wheat was lower at 2.00-2.10. Oats were lower at 2.00-2.10. Hay was lower at 2.00-2.10. Straw was lower at 2.00-2.10. Timber was lower at 2.00-2.10. Lumber was lower at 2.00-2.10. Brick was lower at 2.00-2.10. Cement was lower at 2.00-2.10. Glass was lower at 2.00-2.10. Paper was lower at 2.00-2.10. Textiles were lower at 2.00-2.10. Metals were lower at 2.00-2.10. Minerals were lower at 2.00-2.10. Fuels were lower at 2.00-2.10. Miscellaneous were lower at 2.00-2.10.

**DAIRY AND POULTRY PRODUCTS:** Butter prices were lower, with sales ranging from 250-300 tons. Eggs were lower at 2.00-2.10. Poultry was lower at 2.00-2.10. Milk was lower at 2.00-2.10. Cream was lower at 2.00-2.10. Cheese was lower at 2.00-2.10. Meat was lower at 2.00-2.10. Fish was lower at 2.00-2.10. Seafood was lower at 2.00-2.10. Game was lower at 2.00-2.10. Wildfowl was lower at 2.00-2.10. Domestic fowl was lower at 2.00-2.10. Pigeons were lower at 2.00-2.10. Rabbits were lower at 2.00-2.10. Squirrels were lower at 2.00-2.10. Muskrats were lower at 2.00-2.10. Beavers were lower at 2.00-2.10. Otters were lower at 2.00-2.10. Weasels were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats were lower at 2.00-2.10. Martens were lower at 2.00-2.10. Fishers were lower at 2.00-2.10. Minks were lower at 2.00-2.10. Skunks were lower at 2.00-2.10. Badgers were lower at 2.00-2.10. Foxes were lower at 2.00-2.10. Wolves were lower at 2.00-2.10. Coyotes were lower at 2.00-2.10. Bobcats








**Michigan**  
**ulars regarding your camp**  
**mobile.**

**Detroit,**  
Please mail me full part  
plan for owning an auto  
**Name** \_\_\_\_\_  
**Street** \_\_\_\_\_  
**Town** \_\_\_\_\_  
**IT IS EASY TO OWN A C**

=====









283,167,007.



**"Now, Stay Here Until I Return."**

"Oh, don't bully me!" she protested irritably, trying to control the quaver in her voice. "I won't endure it. Please—"

He looked down at her in silence for a moment, his brown hair in perplexity. Then he turned and went out, settling up the improvised door behind him.

lying motionless in the comparative silence of the night, a prey to rising fears, and, as she lay there, she recalled her shocking fall. However, she faced the fact of her loss bravely. Suffered from all the agonies of recollection with that tranquil

A case for her physical welfare has resulted from clean water. But is something else still held from everyone else, even to the extent of making

It was not to be broken? What might and happen? The fears accumulating grew into feverish terror. She staggered to her feet, and looked toward the door at the door.

The door was barricaded on the outside!

As a aged beast, halfmad with sorrow and impotence, she staggered on and beat the little but hard door.

With a rush of hysterical  
she flung herself upon the  
"Hug!" she cried, dis-  
"Oh, Hug! Hug!"

The sun had moved round behind the RH and the hot seemed dark and

expressive when, suddenly, her eyes  
opened. She started up in some  
alarm. Surely it was not night, and  
"soft" still at least? However auto-  
matically and distasteful any comparison.

---

ered her bewildered brain. She recalled in his grasp, gazing up into his face dazedly. Amid the confusion of her mind his extreme pallor struck her forcibly. His eyes pierced her like flaming steel.

"Hada't you enough sense to realize this possibility?"

Now was the time to assert herself she hesitated; searched vainly for a retort, squared her mouth, closed it again. In her weak state, she

Healing was instantaneous, also there  
by turned her head and eyes toward  
"I was into healing for you," she  
continued, "and by

He said he had been told a man, on a street, with some bad help, had been shot and gone. He stood for some time, and thought, but was unable to go far into the room with eyes so heavily closed with anger, but went.

"If I had been too different then."

For a moment he stood silent, scanning the vicinity of the palm groves, perceiving no signs of the native, he turned with her toward the hut.

luggage," he said, turning the subject of her thoughts abruptly. "I brought some of our things across, and left the rest on the roof. You were asleep. So I took the wireless up the hill and fixed up the serial."

Staring once more upon the coast,  
 She watched him carry in their suit-  
 cases and—something else.  
 To be continued

General Electric orders for the same three months of 1934 were \$40,007,976, 10 per cent better than for the same period of 1933; the year's orders were \$283,167,007.

